

Supplemental Table 3. Comparison of proportion of CoNS according to gestational age between the 2 phases

GA	Phase I (1998–2007)	Phase II (2008–2017)	P value	OR (95% CI)
EOS				
<26 weeks	1/11 (9.1)	2/5 (40.0)	0.214 ^{b)}	6.667 (0.437–101.732)
26–34 weeks	1/11 (9.1)	2/5 (40.0)	0.214 ^{b)}	6.667 (0.437–101.732)
≥35 weeks	9/11 (81.8)	1/5 (20.0)	0.036 ^{b)}	0.056 (0.004–0.805)
LOS				
<26 weeks	13/114 (11.4)	82/345 (23.8)	0.005 ^{a)}	2.422 (1.292–4.541)
26–34 weeks	61/114 (53.5)	169/345 (49.0)	0.402 ^{a)}	0.834 (0.546–1.275)
≥35 weeks	40/114 (35.1)	94/345 (29.2)	0.110 ^{a)}	0.693 (0.441–1.089)

Values are presented as number (%).

CoNS, coagulase-negative staphylococcus; OR, odds ratio; CI, confidence interval; GA, gestational age; EOS, early-onset sepsis; LOS, late-onset sepsis.

^{a)}Chi-square test. ^{b)}Fisher extract test.