

Supplementary Table 2. Notable exclusions

Study	Reasons for exclusions
Nixon (2019) ¹¹	Adults
Sawe (2016) ²	
Choi (2018) ³	
Enright (2010) ⁴	<50 patients
Kwon (2015) ⁵	
Park (2016) ⁶	
Babaie (2018) ⁷	Different target population
Kusumastuti (2021) ⁸	
Mackenzie (2018) ⁹	Not measuring inferior vena cava to aorta ratio
Zhou (2022) ¹⁰	
Vazquez (2017) ¹¹	
Adewumi (2019) ¹²	
Barakat (2018) ¹³	Not using percentage weight difference as the gold standard
Kwon (2016) ¹⁴	
Mazza (2019) ¹⁵	
Koeller (2019) ¹⁶	Same dataset as Modi (2016) ¹⁷
Yunie (2009) ¹⁸	Same dataset as Chen (2007) ¹⁹
Levine (2009) ²⁰	Same dataset as Levine (2010) ²¹
Levine (2010) ²¹	Measures aorta to inferior vena cava ratio
Modi (2016) ¹⁷	

Supplementary References

- Nixon G, Blattner K, Finnie W, Lawrenson R, Kerse N. Use of point-of-care ultrasound for the assessment of intravascular volume in five rural New Zealand hospitals. *Can J Rural Med.* 2019;24(4):109-14.
- Sawe HR, Haefele C, Mfinanga JA, Mwafongo VG, Reynolds TA. Predicting Fluid Responsiveness Using Bedside Ultrasound Measurements of the Inferior Vena Cava and Physician Gestalt in the Emergency Department of an Urban Public Hospital in Sub-Saharan Africa. *PLoS One.* 2016;11(9):e0162772.
- Choi YA, Kwon H, Lee JH, Jung JY, Choi YJ. Comparison of sonographic inferior vena cava and aorta indexes during fluid administered in children. *Am J Emerg Med.* 2018;36(9):1529-33.
- Enright K, Beattie T, Taheri S. Use of a hand-held bladder ultrasound scanner in the assessment of dehydration and monitoring response to treatment in a paediatric emergency department. *Emerg Med J.* 2010;27(10):731-3.
- Kwon H, Lee JH, Kim K, Kwak YH, Kim DK. Measurement of inferior vena cava and aorta with bedside ultrasound to assess degree of dehydration in children. *Critical Ultrasound Journal.* 2015;7(1):A23.
- Park J, Kwon H, Kim D. Ratio of Aorta Area and Inferior Vena Cava Area With Bedside Ultrasound to Assess Degree of Dehydration in Children. *Annals of Emergency Medicine.* 2016;68(4S):S140.
- Babaie S, Behzad A, Mohammadpour M, Reisi M. A Comparison between the Bedside Sonographic Measurements of the Inferior Vena Cava Indices and the Central Venous Pressure While Assessing the Decreased Intravascular Volume in Children. *Adv Biomed Res.* 2018;7:97.
- Kusumastuti NR, Latief A, Pudjiadi AH. Inferior Vena Cava/Abdominal Aorta Ratio as a Guide for Fluid Resuscitation. *J Emerg Trauma Shock.* 2021;14(4):211-5.
- Mackenzie DC, Nasrin S, Atika B, Modi P, Alam NH, Levine AC. Carotid Flow Time Test Performance for the Detection of Dehydration in Children With Diarrhea. *J Ultrasound Med.* 2018;37(6):1397-402.
- Zhou AZ, Green RS, Haines EJ, Vazquez MN, Tay ET, Tsung JW. Interobserver Agreement of Inferior Vena Cava Ultrasound Collapse Duration and Correlated Outcomes in Children With Dehydration. *Pediatr Emerg Care.* 2022;38(1):13-6.
- Vazquez M, Haines E, Tay E, Tsung J. Evaluating Clinical Decisionmaking Using Inferior Vena Cava Ultrasound for IV vs PO Rehydration in Pediatric Emergency Department Patients With Suspected Dehydration. *Annals of Emergency Medicine.* 2017;70(4S):S154.
- Adewumi AA, Braimoh KT, OA MA, Ololu-Zubair HT, Idowu BM. Correlation of sonographic inferior vena cava and aorta diameter ratio with dehydration in Nigerian children. *Niger J Clin Pract.* 2019;22(7):950-6.
- Barakat M, El-Halaby H, Al-Domiatty H. Bedside ultrasonographic evaluation of the degree of dehydration in infants and children with acute gastroenteritis. *Alexandria Journal of Pediatrics.* 2018;31(3):106-11.
- Kwon H, Jung JY, Lee JH, Kwak YH, Kim do K, Jung JH, et al. Sonographic aorta/IVC cross-sectional area index for evaluation of dehydration in children. *Am J Emerg Med.* 2016;34(9):1840-4.
- Mazza G, Romo CM, Torres M, Duffens A, Vyas A, Moran K, et al. Assessment of clinical dehydration using point of care ultrasound for pediatric patients in rural Panama. *World J Emerg Med.* 2019;10(1):46-50.
- Koeller E, Robertson SE, Garbern SC, Monk G, Nasrin S, Alam NH, et al. Ultrasound Adds No Benefit to Clinical Exam for Predicting Dehydration in Children With Acute Diarrhea in a Resource-Limited Setting. *J Ultrasound Med.* 2019;38(3):685-93.
- Modi P, Glavis-Bloom J, Nasrin S, Guy A, Chow A, Dvor N, et al. Accuracy of Inferior Vena Cava Ultrasound for Predicting Dehydration in Children with Acute Diarrhea in Resource-Limited Settings. *PLoS One.* 2016;11(1):e0146859.
- Kim Y. Ultrasound measurement of the inferior vena cava diameter in the assessment of pediatric dehydration: Yale Medicine Thesis Digital Library; 2009.
- Chen L, Kim Y, Santucci KA. Use of ultrasound measurement of the inferior vena cava diameter as an objective tool in the assessment of children with clinical dehydration. *Acad Emerg Med.* 2007;14(10):841-5.
- Levine A, Shah S, Noble V, Epino H. Ultrasound Assessment of Dehydration in Children With Gastroenteritis. *Annals of Emergency Medicine.* 2009;54(3):S91.
- Levine AC, Shah SP, Umulisa I, Munyaneza RB, Dushimiyimana JM, Stegmann K, et al. Ultrasound assessment of severe dehydration in children with diarrhea and vomiting. *Acad Emerg Med.* 2010;17(10):1035-41.