

Supplementary References

1. Equivalent opioid calculator [Internet]. ClinCalc LLC; 2025 [cited 2022 Feb 20]. Available from: <https://clincalc.com/Opioids>.
2. Equivalent benzodiazepine calculator [Internet]. ClinCalc LLC; 2025 [cited 2022 Feb 20]. Available from: <https://clincalc.com/Benzodiazepine>.
3. Blackwood B, Tume LN, Morris KP, Clarke M, McDowell C, Hemming K, et al. Effect of a sedation and ventilator liberation protocol vs usual care on duration of invasive mechanical ventilation in pediatric intensive care units: a randomized clinical trial. *JAMA* 2021;326:401-10.
4. Chomat MR, Said AS, Mann JL, Wallendorf M, Bickhaus A, Figueroa M. Changes in sedation practices in association with delirium screening in infants after cardiopulmonary bypass. *Pediatr Cardiol* 2021;42:1334-40.
5. Curley MA, Wypij D, Watson RS, Grant MJ, Asaro LA, Cheifetz IM, et al. Protocolized sedation vs usual care in pediatric patients mechanically ventilated for acute respiratory failure: a randomized clinical trial. *JAMA* 2015;313:379-89.
6. Deeter KH, King MA, Ridling D, Irby GL, Lynn AM, Zimmerman JJ. Successful implementation of a pediatric sedation protocol for mechanically ventilated patients. *Crit Care Med* 2011;39:683-8.
7. Gaillard-Le Roux B, Liet JM, Bourgoin P, Legrand A, Roze JC, Joram N. Implementation of a nurse-driven sedation protocol in a PICU decreases daily doses of midazolam. *Pediatr Crit Care Med* 2017;18:e9-17.
8. Hanser A, Neunhoffer F, Hayer T, Hofbeck M, Schlensak C, Mustafi M, et al. A nurse-driven analgesia and sedation protocol reduces length of PICU stay and cumulative dose of benzodiazepines after corrective surgery for tetralogy of Fallot. *J Spec Pediatr Nurs* 2020;25:e12291.
9. Jin HS, Yum MS, Kim SL, Shin HY, Lee EH, Ha EJ, et al. The efficacy of the COMFORT scale in assessing optimal sedation in critically ill children requiring mechanical ventilation. *J Korean Med Sci* 2007;22:693-7.
10. Keogh SJ, Long DA, Horn DV. Practice guidelines for sedation and analgesia management of critically ill children: a pilot study evaluating guideline impact and feasibility in the PICU. *BMJ Open* 2015;5:e006428.
11. Kleiber N, de Wildt SN, Cortina G, Clifford M, van Rosmalen J, van Dijk M, et al. A comparative analysis of preemptive versus targeted sedation on cardiovascular stability after high-risk cardiac surgery in infants. *Pediatr Crit Care Med* 2016;17:321-31.
12. Kozin ED, Cummings BM, Rogers DJ, Lin B, Sethi R, Noviski N, et al. Systemwide change of sedation wean protocol following pediatric laryngotracheal reconstruction. *JAMA Otolaryngol Head Neck Surg* 2015;141:27-33.
13. Neunhoffer F, Kumpf M, Renk H, Hanelt M, Berneck N, Bosk A, et al. Nurse-driven pediatric analgesia and sedation protocol reduces withdrawal symptoms in critically ill medical pediatric patients. *Paediatr Anaesth* 2015;25:786-94.

14. Neunhoeffter F, Seitz G, Schmidt A, Renk H, Kumpf M, Fideler F, et al. Analgesia and sedation protocol for mechanically ventilated postsurgical children reduces benzodiazepines and withdrawal symptoms-but not in all patients. *Eur J Pediatr Surg* 2017;27:255-62.
15. Saelim K, Chavananon S, Ruangnapa K, Prasertsan P, Anuntaseree W. Effectiveness of protocolized sedation utilizing the COMFORT-B scale in mechanically ventilated children in a pediatric intensive care unit. *J Pediatr Intensive Care* 2019;8:156-63.
16. Sanchez-Pinto LN, Nelson LP, Lieu P, Koh JY, Rodgers JW, Larson KA, et al. Implementation of a risk-stratified opioid weaning protocol in a pediatric intensive care unit. *J Crit Care* 2018;43:214-9.
17. Solodiuk JC, Greco CD, O'Donnell KA, Morrill DR, Curley MAQ. Effect of a sedation weaning protocol on safety and medication use among hospitalized children post critical illness. *J Pediatr Nurs* 2019;49:18-23.
18. Tiacharoen D, Lertbunrian R, Veawpanich J, Suppalarkbunlue N, Anantasit N. Protocolized sedative weaning vs usual care in pediatric critically ill patients: a pilot randomized controlled trial. *Indian J Crit Care Med* 2020;24:451-8.
19. Verlaet CW, Heesen GP, Vet NJ, de Hoog M, van der Hoeven JG, Kox M, et al. Randomized controlled trial of daily interruption of sedatives in critically ill children. *Paediatr Anaesth* 2014;24:151-6.
20. Wilson AK, Ragsdale CE, Sehgal I, Vaughn M, Padilla-Tolentino E, Barczyk AN, et al. Exposure-based methadone and lorazepam weaning protocol reduces wean length in children. *J Pediatr Pharmacol Ther* 2021;26:42-9.
21. Achuff BJ, Lemming K, Causey JC, Sembera KA, Checchia PA, Heinle JS, et al. Opioid weaning protocol using morphine compared with nonprotocolized methadone associated with decreased dose and duration of opioid after norwood procedure. *Pediatr Crit Care Med* 2022;23:361-70.
22. Alfraj, A., et al., Effect of target-driven sedation protocol to ventilator liberation in pediatric intensive care unit: pre- and postimplementation single-center study. *J Pediatr Intensive Care* 2023. <https://doi.org/10.1055/s-0043-1774306>.
23. Dokken M, Rustøen T, Diep LM, Fagermoen FE, Huse RI, Egerod I, et al. Implementation of an algorithm for tapering analgosedation reduces iatrogenic withdrawal syndrome in pediatric intensive care. *Acta Anaesthesiol Scand* 2023;67:1229-38.
24. King CE, Wood DN, Koo J, Cutler AB, Vesel TP. Sedation weaning initiative targeting methadone exposure: single center improvements in withdrawal symptoms and hospital length of stay for pediatric cardiac critical care. *Pediatr Crit Care Med* 2023;24:e332-41.
25. Leu GR, Scott AR. A standardized care pathway following mandibular distraction in infants less than 3 months of age. *Otolaryngol Head Neck Surg* 2019;161:870-6.

26. Loberger JM, Waddell KC, Prabhakaran P, Jones RM, Lawrence MV, Bittles LA, et al. Pediatric ventilation liberation: bundled extubation readiness and analgosedation pathways decrease mechanical ventilation duration and benzodiazepine exposure. *Respir Care* 2022;67:1385-95.
27. Shildt N, Traube C, Dealmeida M, Dave I, Gillespie S, Moore W, et al. "Difficult to sedate": successful implementation of a benzodiazepine-sparing analgosedation-protocol in mechanically ventilated children. *Children (Basel)* 2021;8:348.
28. Taffarel P, Widmer J, Fiore Á, Rodríguez AP, Meregalli C, Jorro Barón F. Impact of the implementation of a sedation and analgesia protocol in a pediatric intensive care unit. *Arch Argent Pediatr* 2023;121:e202202806. English, Spanish.
29. Ambuel B, Hamlett KW, Marx CM, Blumer JL. Assessing distress in pediatric intensive care environments: the COMFORT scale. *J Pediatr Psychol* 1992;17:95-109.
30. van Dijk M, Peters JW, van Deventer P, Tibboel D. The COMFORT Behavior Scale: a tool for assessing pain and sedation in infants. *Am J Nurs* 2005;105:33-6.
31. Curley MA, Harris SK, Fraser KA, Johnson RA, Arnold JH. State Behavioral Scale: a sedation assessment instrument for infants and young children supported on mechanical ventilation. *Pediatr Crit Care Med* 2006;7:107-14.
32. Traube C, Silver G, Kearney J, Patel A, Atkinson TM, Yoon MJ, et al. Cornell Assessment of Pediatric Delirium: a valid, rapid, observational tool for screening delirium in the PICU*. *Crit Care Med* 2014;42:656-63.
33. Franck LS, Harris SK, Soetenga DJ, Amling JK, Curley MA. The Withdrawal Assessment Tool-1 (WAT-1): an assessment instrument for monitoring opioid and benzodiazepine withdrawal symptoms in pediatric patients. *Pediatr Crit Care Med* 2008;9:573-80.
34. Ista E, van Dijk M, de Hoog M, Tibboel D, Duivenvoorden HJ. Construction of the Sophia Observation withdrawal Symptoms-scale (SOS) for critically ill children. *Intensive Care Med* 2009;35:1075-81.
35. Ramelet AS. Assessment of pain and agitation in critically ill infants. *Aust Crit Care* 1999;12:92-6.
36. Ramelet AS, Rees N, McDonald S, Bulsara M, Abu-Saad HH. Development and preliminary psychometric testing of the Multidimensional Assessment of Pain Scale: MAPS. *Paediatr Anaesth* 2007;17:333-40.
37. Ramelet AS, Rees NW, McDonald S, Bulsara MK, Huijter Abu-Saad H. Clinical validation of the multidimensional assessment of pain scale. *Paediatr Anaesth* 2007;17:1156-65.
38. Franck L, Vilardi J. Assessment and management of opioid withdrawal in ill neonates. *Neonatal Netw* 1995;14:39-48.
39. Franck LS, Vilardi J, Durand D, Powers R. Opioid withdrawal in neonates after continuous infusions of morphine or fentanyl during extracorporeal membrane oxygenation. *Am J Crit Care* 1998;7:364-9.

40. Thottam PJ, Gilliland T, Ettinger N, Baijal R, Mehta D. Outcomes using a postoperative protocol in pediatric single-stage laryngotracheal reconstruction. *Ann Otol Rhinol Laryngol* 2021;130:861-7.